This Page Is Inserted by IFW Operations and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.

NO DRAWINGS

Prof. (1000)

proceedings of the conditional

ر(نالئ 🗒

Profit to the

NEW ZEALAND

PATENTS ACT 1953

No

Date

COMPLETE SPECIFICATION

IMPROVED PERSONAL SKIN CARE PRODUCTS CONTAINING DIMETHYE DIALLYE AMMORTHM CHEORIDE/ACTYLIC ACID-TYPE FOLYMERS.

HAWE. CALGON CORPORATION, a corporation duly organized and existing under the laws of the State of Delaware, United States of America of Houte 60-(amphell's Run Road, Poblisson fownship, State of Lorentz Lania, United States of America,

hereby declare the invention for which 17/ we pray that a patent may be granted to medus, and the method by which it is to be performed, to be particularly described in and by the following statement:-

C-1442

TITLE OF THE INVENTION

"IMPROVED PERSONAL SKIN CARE PRODUCTS CONTAINING DIMETHYL DIALLYL AMMONIUM CHLORIDE/ACRYLIC ACID-TYPE POLYMERS"

BACKGROUND OF THE INVENTION

The instant invention relates to improved personal skin care products which contain at least one dimethyl diallyl ammonium chloride (DMDAAC)/ acrylic acid-type polymer. These polymers impart a soft, silky feel to the skin when present at low concentrations and are compatible with anionic systems.

Numerous references disclose cosmetic and personal care compositions such as shampoos, antiperspirant formulations, anti-dandruff rinse conditioners, etc., which contain anionic and/or cationic polymers, an active agent, surfactants, emollients and other additives and preservatives commonly employed in the industry. Pertinent references which relate to

- 2 - 144.

DMDAAC/acrylic acid polymers or which disclose cosmetic and personal care formulations containing anionic and cationic polymers include:

1. US Patent 3,761,417, which is directed to detergent compositions containing particle deposition enhancing agents. More particularly, this patent discloses detergent and personal use toilet detergent bars containing water-soluble particles such as antimicrobial agents, organic surfactants and cationic polymers. Surfactants are an essential ingredient of these compositions, and DMDAAC is mentioned as a possible cationic polymer.

2. US Patent 3,769,398, which discloses non-ronic hair shampoo formulations containing an active ingredient such as a betaine, sulfo betaine, amine oxide or mixture thereof, a water soluble polymer such as a polyethyleneimine-ethylene oxide or

propylene oxide polymer or a propoxylated polyethyleneimine.

3. US Patent 4,329,335, which describes an amphoteric, nonionic anti-dandruff shampoo containing an active agent (1-imidazaly1-1-) (chlorophenoxy-3, 3-dimethylbutane-2-one) and amphoteric surfactants. DMDAAC is disclosed as a preferred quaternized ammonium compound in this patent.

30

10

i 5

4. Published European Patent Application No. 74,819 which discloses an anti-dandruff cream ringe conditioner containing zinc pyrithione, glucus guar gum, hydroxyethyl cellulose and a homopolymer of DMDAAC or a copolymer of DMDAAC and acrylamide.

5. US Patent 3,996,146, which discloses a shampoof formulation comprising from 0.05 to about 2.5%, by weight, of a cationic resin including quaternary polymers derived from dimethyl dially ammonium salts.

10

15

- 6. US Patent 4,040,984, which discloses polymers useful for preparing electroconductive paper which comprise quaternary diallyl dialkyl ammonium monomers and acrylic acid.
- 7. US Patent 3,912,808, which discloses a composition and method for waving or straightening hair using an aqueous solution of a reducing agent and a water soluble secondary or tertiary amine polymer or a polymer of diallyl amine or a quaternary polymer of diallyl dialkyl ammonium salts. This patent also discloses the use of dialkyl ammonium polymers which contain acrylamide or diacetone acrylamide. The use of dimethyl diallyl ammonium chloride/acrylic acid polymers is not disclosed or suggested.
- 30 8. US Patent 4,027,008, which discloses hair treating compositions which contain a water soluble secondary or tertiary amine polymer or a polymer of diallyl amine or a quaternary polymer of

dially! dialky! ammonium salts. This patent in not disclose or suggest the use of DMDAAF acryl: acid polymers, and states that many winely well products for treating hair contain animals surfactants which may inactivate cationic additives.

9. US Patent 3,986,825, which discloses the use of dialkyldiallyl ammonium polymers in cosmetic products, including copolymers of a dialkyl diallyl ammonium monomer and acrylamide or diacetone acrylamide. However, polymers containing these cationic moieties and acrylic acid are not disclosed or suggested.

15

In summary, though dialkyl diallyl ammonium polymers are widely used in cosmetic applications, the use of polymers containing a diallyl dialkyl ammonium monomer and acrylic acid is not known or suggested in the art. These polymers provide an increased level of cationic activity with unexpected compatibility in anionic systems.

DETAILED DESCRIPTION OF THE INVENTION

25

20

The instant invention is directed to a method forimproving the feel of a personal skin care product, comprising adding to said product an effective amount of a polymer comprising:

30

a) substantially 60 to 99%, pased on this polyner weight, of a quaternary diallyl dialkyl ammonium monomer, wherein alkyl groups are independently selected from alkyl groups of l



to 18 carbon atoms, preferably organization, and wherein said quaternary dially indiality, ammonium monomer's counterion in solution from the arrosp consisting of conjugate taken

of acids having an ionization constant greater than 10⁻¹³, more preferably, selected from the group consisting of fluoride, bromide, chloride, hydroxide, nitrate, acetate, hydrogen sulfate, and primary phosphates; and

10

15

20

25

30

polymer weight, not an automorphism to prove the polymer weight, not an automorphism of acrylic acid and methacrylic acid; wherein the weight average molecular weight of said polymer tander from materially 11,000 to our tentralie in a determined in mediance increases.

The instant invention is also directed to personal skin care products which contain the above described polymer.

As used herein, the phrase "personal ckin care product" includes, but is not limited to, nano and face lotions, soaps and creams, suntan lotions, bubble baths, shaving creams, antiperupirants and deodorants. The instant polymers impart a soft and silky feel to the skin when products containing them are used. The term "feel", at used herein, refers to a subjective measure of the smoothness and silkiness that a personal skin care product imparts to the skin when applied to the skin.

1-144%

Personal skin care products generally comprise an active agent, such as a detergent or surfactant, conditioner, emollient, antimicrotial, antiperspirant

and moisturizing agent. For example, active agents may include antimicrobial agents such as trichlorocarban (3,4,4'-trichloro carbanilide), triclosan (2,4,4'-trichloro carbonilide), triclosan (2,4,4'-trichloro-2'-hydroxy diphenyl ether), benzalkonium chloride, zinc phenosulfonate, zinc ricinoleate and the like.

Commonly used anti-perspirants include aluminum zirconium complex, aluminum chlorohydrate and the like.

Representative emollients, humectants and moisturizing agents include C₁₂-C₁₅ alcohol benzoates, sorbitol, glycerin, propylene glycol (PEG), lanolin, vegetable oils, mineral oils, isopropyl myristate, aloe vera, jojoba oil and the like.

20

25

30

ì 0

Other cosmetically acceptable excipients that a personal care product formulation may contain are thickening agents, buffering agents and preservatives. Suitable water soluble preservatives are sodium bisulfite, sodium thiosulfate, ascorbate, benzalkonium chloride, glydant chlorobutanol, thimerosal, phenylmercuric borate, Dowicil 200 parabens, Tektamer 38(1,2-dibromo-2,4-dicyanobutane), benzyl alcohol, phenylethanol and the like. Suitable thickening agents are Cab-O-Sil M5 made by Cabot Corporation, sodium stearate, magnesium aluminum silicate, hydroxyethyl cellulose and the like. These agents may be present in amounts of from 0.05 to 50% by weight and preferably 1 to 5%. Suitable water

soluble buffering agents are alkali or alkali earen carbonates, phosphates, bicarbonates, citrates, borates, acetates, acid anhydrides, succinates and the

like, tuch as sodium phosphate, citrate, borate, acetate, bicarbonate and carbonate. These agents may be present in an amount sufficient to maintain some optimum pH of the system in the range 2 to 9. As such, the buffering agent can be as much as 20%, on a weight basis, of the total composition. Additional active agents are fully described in U.S. Fatent 3,986,825, which is incorporated herein by reference.

The instant polymers, in addition to imparting improved feel to skin, may enhance dispersion and improve the efficacy of functional ingredients in personal skin care products.

10

15

20

25

As used herein, the term "effective amount" refers to that amount of polymer required to improve the (sel of the personal skin care product to which it is added. Generally, the instant polymers are added at a dosage ranging from substantially 0.1 amount of the composition to substantially 1. active polymer solids. hose there the trail weight of the product to which the polymer is added. Preferably, the dosage ranges from about 0.2 to about 3%, based on the total weight of the composition being treated and most preferably from about 0.5% to about 2.5%, based on the total weight of the composition being treated.

The quaternary diallyl dialkyl ammonium monomer comprises from substantially 60 to substantially 60 to substantially 60 to substantially 6 to substantially 1 to substantially 60, based on total polymer.

The state of the s

weight. Preferably, the quaternary: animon weight ratio is from about 95:5 to about 75:25, based on the total weight of polymer. Thus, in the polymers of the present invention, the cationic molety of the polymer is predominant while the anionic molety of the polymer is minor. Additionally, other moleties may be present in the instant polymers.

An especially suitable polymer is that where the cationic portion is dimethyldiallyl ammonium chloride (DEDAAC) (DMDAAC) or diethyldiallyl ammonium chloride (DEDAAC) and where anionic portion is acrylic acid.

Preferably, the DMDAAC/DEDAAC:acrylic acid weight ratio weight ratio ranges from about 99:1 to about 60:40, most preferably from about 95:5 to 75:25, based on total polymer weight.

The polymers of the instant invention may have any molecular weight ranging from substantially 10,000,000, with the preferred molecular weight ranging from about 200,000 to about 5,000,000. The most preferred viscosity for the instant polymers ranges from about 4,000 to about 10,000 cps, as determined using a Brookfield LVF No. 4 spindle at 60 rpm. These polymers may be prepared using any conventional free radical polymerization technique, such as the technique disclosed by Butler and Angelo, "Journal of American Chemical Society," Vol. 79, p. 3128 (1957) or the technique suggested in C.S. Re sue Patent No. 28,543. These references are incorporated by reference into this specification.

20

25

30



EXAMPLE 1

10

15

The following example is not intended to limit to

scope of this invention in any way.

This example demonstrates the use of the instant polymers in anionic personal skin care product formulations.

PEARLESCENT LIQUID SOAP

A liquid soap formulation was prepared using the following ingredients in accordance with the following instructions to demonstrate the excellent compatibility of the instant polymers with anionic surfactants. Compatibility of Merquat 280, which has a net cationic charge density, with an anionic system is surprising and unexpected.

20	Ingredients		<u>* * * * * * </u>
	F.	Deionized Water	€ <u>1</u> . 9 (
		Sodium Lauryl Sulfate	25.00
		Sodium C ₁₄₅ 16 Olefin Sulfo	onate ¹ 10.00 %
25		Cerasynt M	0.40
		Cocamide DEA ³	2.60
		Cocamidopropyl Betaine ⁴	8.00
30	В	MERQUAT 280 ⁵	2.00
	C	Fragrance	0.50
		Tetrasodium EDTA ⁶	0.20
	D	Citric Acid	to pH 6.00
35		Sodium Chloride	to viscosity 2800 cps

Preparation Instructions

10

15

Heat water to 75°C. With moderate agricultation, the the ingredients listed in Part A in the order states proceeding after each addition is uniform. Beauty cooling and add MERQUAT 200 at 50°C. Continue mixing and cooling and add Part C at 45°C. Adjust pR to 6.0 with outric acid. Add sodium chloride to increase viscolity to 2800 cps. Optionally, fragrances and dyes can be added.

- 1. Sodium laureth sulfate is C_{14-16} olefin sulfonate sulfate is an anionic surfactant.
- Cerasynt is ethylene glycol monostearate, available from Van Dyk & Co., inc.
- Cocamide DEA is coconut diethanolamine, available from Van Dyk and Co., Inc.
- 4. Cocamidopropyl betaine is a coconut asidepropyl betaine.
- 5. Merquat 280 is an 80/20 w/w polymer of DMDAAC and acrylic acid having a viscosity (Brookfield DVF #4 spindle 030 rpm) of 4,000-10,000 cp., and a weight average molecular weight of approximately 1,300,000, as determined by gel permeation chromatography, commercially available from Galace Corporation.
 - 5. Tetrasodium EDTA is the sodium salt of ethlenediamine tetraacetic acid.

WHAT WE CLAIM IS:

- l. A method for improving the feel of a personal skin care product, comprising adding to said product an effective amount of a polymer comprising:
 - a. substantially 60 to substantially 99%, by weight of said polymer, of a quaternary diallyl dialkyl ammonium monomer, wherein alkyl groups are independently selected from alkyl groups of 1 to 18 carbon atoms and wherein said quaternary diallyl dialkyl ammonium monomer's counterion is selected from the group consisting of conjugate bases of acids having an ionization constant greater than 10-13; and
 - b. substantially 1 to substantially 40%, by weight of said polymer, of an anionic monomer selected from the group consisting of acrylic acid and methacrylic acid; wherein the weight average molecular weight of said polymer ranges from substantially 50,000 to 10,000,000, as determined by gel permeation chromatography.
 - 2. The method of Claim 1, wherein said alkyl group of a) is C_{1-4} .
- 3. The method of Claim 1, wherein a) is selected from the group consisting of dimethyldiallyl ammonium chloride and diethyldiallyl ammonium chloride and b) is acrylic acid.

- 4. The method of any one of Claims 1 to 3, wherein said effective amount ranges from substantially 0.1 to substantially 5%, by weight, of said product.
- 5. The method of any one of Claims 1 to 4, wherein said product comprises an active agent and a surfactant.
- 6. The method of Claim 5, wherein said surfactant is anionic.
- 7. The method of any one of Claims 1 to 6, wherein said counterion is selected from the group consisting of fluoride, chloride, bromide, hydroxide, nitrate, acetate, hydrogen sulfate and primary phosphates.
- 15 8. A personal skin care product which comprises:
 - a. an active agent;
 - b. an anionic surfactant;
- 20 c. a copolymer comprising:
 - 1. substantially 60 to substantially 99%, by weight of said polymer, of a quaternary diallyl dialkyl ammonium monomer, wherein alkyl groups are independently selected from alkyl groups of 1 to 18 carbon atoms and wherein said quaternary diallyl dialkyl ammonium monomer's counterion is selected from the group consisting of conjugate bases of acids having an ionization constant greater than 10-13; and

substantially 1 to substantially 40%, by weight of said polymer, of a monomer selected from group consisting of acrylic acid and methacrylic acid; wherein said polymer has a weight average molecular weight ranging from substantially 50,000 to substantially 10,000,000, as determined by gel permeation chromatography.

30

25



9. The product of Claim 8 wherein 1. is selected from the group consisting of dimethyldiallyl ammonium chloride and diethyldiallyl ammonium chloride and 2. is acrylic acid.

28 august 10 90 Greta Gottleb.

